

1. When you get home from surgery

- Start using all 3 medication drops on the regular dosing schedule.
ANTIBIOTIC, **NSAID**, and **PREDNISOLONE** are all 3 times a day.
- Do not rub your eye or press on it.
- Sleep with the shield tonight.

ANTIBIOTIC



2. The day after the surgery

- You do not need to wear the shield any more.
- You can resume normal activities.
- Bending and lifting are OK as long as there is no discomfort.
- Showering or bathing are OK but avoid getting water in your eye.
- Do not wear makeup for 1 week.
- Bring all of your drops with you to each appointment

NSAID



3. Postoperative eye drops

- ANTIBIOTIC** 1 drop 3 times a day for 10 days.
- NSAID** 1 drop 3 times a day for 1 month.
- PREDNISOLONE** 1 drop 3 times a day for 1 month.
Shake this one medication well before using.
- Wait a few minutes between each drop.



4. Normal postoperative complaints

- Your eye may be red.
- Your vision may be blurred.
- Your pupil may be large and dilated.
- Your eye may feel scratchy, teary, and light sensitive.
- You may see floaters or black spots.

PREDNISOLONE



- ## 5. Please call the office immediately if there is a serious problem.
- Increased redness, pain, swelling, or blurred vision requires an immediate phone call.