

1. When you get home from surgery

- a. Start using all 3 medication drops on the regular dosing schedule.
ANTIBIOTIC, **NSAID**, and **PREDNISOLONE** are all 3 times a day.
- b. Do not rub your eye or press on it.
- c. Sleep with the shield tonight.

ANTIBIOTIC



2. The day after the surgery

- a. You do not need to wear the shield any more.
- b. You can resume normal activities.
- c. Bending and lifting are OK as long as there is no discomfort.
- d. Showering or bathing are OK but avoid getting water in your eye.
- e. Do not wear makeup for 1 week.
- f. Bring all of your drops with you to each appointment

NSAID



3. Postoperative eye drops

- a. **ANTIBIOTIC** 1 drop 3 times a day for 10 days.
- b. **NSAID** 1 drop 3 times a day for 1 month.
- c. **PREDNISOLONE** 1 drop 3 times a day for 1 month.
Shake this one medication well before using.
- d. Wait a few minutes between each drop.



4. Normal postoperative complaints

- a. Your eye may be red.
- b. Your vision may be blurred.
- c. Your pupil may be large and dilated.
- d. Your eye may feel scratchy, teary, and light sensitive.
- e. You may see floaters or black spots.

PREDNISOLONE



5. Please call the office immediately if there is a serious problem. Increased redness, pain, swelling, or blurred vision requires an immediate phone call.